

**Synonym**

Cadherin-17,CDH17,HPT-1,LI-cadherin

**Source**

Biotinylated Human Cadherin-17 Protein, Fc,Avitag(CA7-H82F9) is expressed from human 293 cells (HEK293). It contains AA Gln 23 - Met 787 (Accession # [Q12864-1](#)).

Predicted N-terminus: Gln 23

**Molecular Characterization**

Cadherin-17(Gln 23 - Met 787) Q12864-1	Fc(Pro 100 - Lys 330) P01857	Avi
---	---------------------------------	-----

This protein carries a human IgG1 Fc tag at the C-terminus, followed by an Avi tag (Avitag™)

The protein has a calculated MW of 88.6 kDa. The protein migrates as 43-140 kDa when calibrated against [Star Ribbon Pre-stained Protein Marker](#) under reducing (R) condition (SDS-PAGE) due to glycosylation.

**Labeling**

*Biotinylation of this product is performed using Avitag™ technology. Briefly, the single lysine residue in the Avitag is enzymatically labeled with biotin.*

**Protein Ratio**

Passed as determined by the HABA assay / binding ELISA.

**Endotoxin**

Less than 1.0 EU per µg by the LAL method.

**Purity**

>90% as determined by SDS-PAGE.

>90% as determined by SEC-MALS.

**Formulation**

Lyophilized from 0.22 µm filtered solution in 20 mM Tris, 150 mM NaCl, pH8.0 with trehalose as protectant.

Contact us for customized product form or formulation.

**Reconstitution**

Please see Certificate of Analysis for specific instructions.

*For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.*

**Storage**


For long term storage, the product should be stored at lyophilized state at -20°C or lower.

*Please avoid repeated freeze-thaw cycles.*

This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.


**SDS-PAGE**



Biotinylated Human Cadherin-17 Protein, Fc,Avitag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 90% (With [Star Ribbon Pre-stained Protein Marker](#)).


**Bioactivity-ELISA**

**SEC-MALS**



The purity of Biotinylated Human Cadherin-17 Protein, Fc,Avitag (Cat. No. CA7-H82F9) is more than 90% and the molecular weight of this protein is around 275-315 kDa verified by SEC-MALS.

[Report](#)



Immobilized Anti-CDH17 antibody at 0.2% can bind Biotinylated Human Cadherin-17 Protein, Fc,Avitag (Cat. No. CA7-H82F9) with a linear range of 0.039-1.25 µg/mL (QC tested).

## Background

Cadherin-17, also known as liver-intestine (LI) Cadherin, belongs to the cadherin family of calcium-dependent cell adhesion molecules. In vivo studies showed CDH17 knockout resulted in apoptotic PC tumor death through activating caspase-3 activity. Taken together, CDH17 functions as an oncogenic molecule critical to PC growth by regulating tumor apoptosis signaling pathways and CDH17 could be targeted to develop an anti-PC therapeutic approach.

## Clinical and Translational Updates

Please contact us via [TechSupport@acrobiosystems.com](mailto:TechSupport@acrobiosystems.com) if you have any question on this product.