



Synonym

CD3E & CD3D, CD3 delta & CD3 epsilon

Source

Cynomolgus CD3E&CD3D Heterodimer Protein, Fc,His Tag&Fc,Flag Tag(CDD-C52W9) is expressed from human 293 cells (HEK293). It contains AA Gln 22 - Asp 117 (CD3E) & Phe 22 - Ala 105 (CD3D) (Accession # [Q95LI5-1](#) (CD3E) & [Q95LI8-1](#) (CD3D)).

Predicted N-terminus: Gln 22 (CD3E) & Phe 22 (CD3D)

Molecular Characterization

CD3E (Gln 22 - Asp 117) Q95LI5-1	Fc(Pro 100 - Lys 330) P01857	Poly-his
CD3D (Phe 22 - Ala 105) Q95LI8-1	Fc(Pro 100 - Lys 330) P01857	Flag

Cynomolgus CD3E&CD3D Heterodimer Protein, Fc,His Tag&Fc,Flag Tag is produced by co-expression of CD3E and CD3D, has a calculated MW of 42.5 kDa (CD3E) and 40.9 kDa (CD3D). Subunit CD3E is fused with a human IgG1 Fc tag and a polyhistidine tag at the C-terminus and subunit CD3D is fused with a human IgG1 Fc tag and a flag tag at the C-terminus. The predicted N-terminus is Gln 22 (CD3E) & Phe 22 (CD3D). The reducing (R) protein migrates as 46-50 kDa and 55-65 kDa due to glycosylation.

Endotoxin

Less than 1.0 EU per µg by the LAL method.

Purity

>90% as determined by SDS-PAGE.

>90% as determined by SEC-MALS.

Formulation

Lyophilized from 0.22 µm filtered solution in PBS, pH7.4 with trehalose as protectant.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.

Storage


For long term storage, the product should be stored at lyophilized state at -20°C or lower.

Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.


SDS-PAGE



Cynomolgus CD3E&CD3D Heterodimer Protein, Fc,His Tag&Fc,Flag Tag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 90%.

Bioactivity-ELISA

SEC-MALS



The purity of Cynomolgus CD3E&CD3D Heterodimer Protein, Fc,His Tag&Fc,Flag Tag (Cat. No. CDD-C52W9) is more than 90% and the molecular weight of this protein is around 90-120 kDa verified by SEC-MALS.

[Report](#)

Discounts, Gifts,
and more!



Cynomolgus CD3 epsilon&CD3 delta Heterodimer Protein, Fc,His Tag&Fc,Flag Tag (MALS verified)

Catalog # CDD-C52W9




BIOSYSTEMS
Acro

Surprise Inside!


Cynomolgus CD3E&CD3D Heterodimer Protein, Fc,His Tag&Fc,Flag Tag ELISA

0.1 µg of Bispecific CD3×BCMA T cell-engaging Antibody per well



Immobilized Bispecific CD3×BCMA T cell-engaging Antibody at 1 µg/mL (100 µL/well) can bind Cynomolgus CD3E&CD3D Heterodimer Protein, Fc,His Tag&Fc,Flag Tag (Cat. No. CDD-C52W9) with a linear range of 1-20 ng/mL (QC tested).

Bioactivity-BLI



Loaded Cynomolgus CD3E&CD3D Heterodimer Protein, Fc,His Tag&Fc,Flag Tag (Cat. No. CDD-C52W9) on AHC Biosensor, can bind Bispecific CD3×BCMA T cell-engaging Antibody with an affinity constant of 1.77 nM as determined in BLI assay (ForteBio Octet Red96e) (Routinely tested).

Background

T-cell surface glycoprotein CD3 delta & CD3 epsilon chain, also known as CD3D & CD3E or CD3D&CD3E respectively, are single-pass type I membrane proteins. CD3D, together with CD3- epsilon(CD3E) , CD3-gamma and CD3-zeta, and the T-cell receptor alpha/beta and gamma/delta heterodimers, forms the T cell receptor-CD3 complex. T cell receptor-CD3 complex plays an important role in coupling antigen recognition to several intracellular signal-transduction pathways.

Clinical and Translational Updates

Discounts, Gifts,
and more!



➤ www.acrobiosystems.com

5/8/2024