

HRP conjugated Monoclonal Anti-DM-1&DM-4 Antibody, Mouse IgG1

Catalog # DM1-PLY73



BIOSYSTEMS
Acro

Source

HRP conjugated Monoclonal Anti-DM-1&DM-4 Antibody, Mouse IgG1 is a Mouse monoclonal antibody produced from a hybridoma created by fusing SP2/0 myeloma and Mouse B-lymphocytes.

Species

Mouse

Isotype

Mouse IgG1 | Mouse Kappa

Conjugate

HRP-Conjugated

Antibody Type

Hybridoma Monoclonal

Reactivity

Chemical

Immunogen

DM-1.

Specificity

This product is a specific antibody specifically reacts with DM-1 and DM-4.

Application

Application	Recommended Usage
ELISA	1-500 ng/mL

Purity

>90% as determined by SDS-PAGE.

Purification

Protein A purified/ Protein G purified

Formulation

Lyophilized from 0.22 µm filtered solution in PBS, pH7.4 with trehalose as protectant.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.

Storage


For long term storage, the product should be stored at lyophilized state at -20°C or lower.

Please protect from light and avoid repeated freeze-thaw cycles.

This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.

SDS-PAGE



HRP conjugated Monoclonal Anti-DM-1&DM-4 Antibody, Mouse IgG1 on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie

Discounts, Gifts,
and more!



➤ www.acrobiosystems.com

9/2/2024

HRP conjugated Monoclonal Anti-DM-1&DM-4 Antibody,Mouse IgG1


Catalog # DM1-PLY73



BIOSYSTEMS
Acro

Blue. The purity of the protein is greater than 90% (With [Star Ribbon Pre-stained Protein Marker](#)).

Bioactivity-ELISA



Immobilized Trastuzumab-DM1 (T-DM1) at 2 µg/mL (100 µL/well) can bind HRP conjugated Monoclonal Anti-DM-1&DM-4 Antibody,Mouse IgG1 (Cat. No. DM1-PLY73) with a linear range of 0.24-15.63 ng/mL (QC tested).

Background

Mertansine (DM-1) is a tubulin inhibitor that binds to the ends of microtubules and inhibits microtubule dynamics. DM-1(Mertansine) has antitumor activity and functions as a regulator of tubulin. It is an alpha-amino acid ester, a carbamate, an epoxide, an organic heterocyclic tetracyclic compound, an organochlorine compound, a mercaptan, and a maydenin alkaloid. DM-1, derived from Mydenin, is a cytotoxic component of antibody-drug conjugations that produce antibody-drug conjugations via a sulfhydryl group splice with SPP (n-succinimide 4- (2-pyridyl dithio)) or SMCC (4- (3-mercapto-2, 5-dioxy-1 pyrrolidyl) -cyclohexanic acid) splice.

Clinical and Translational Updates

Discounts, Gifts,
and more!



➤ www.acrobiosystems.com

9/2/2024