

# **Synonym**

DEL1,EDIL3

#### Source

Human EDIL3, Fc Tag(ED3-H5259) is expressed from human 293 cells (HEK293). It contains AA Asp 24 - Glu 480 (Accession # O43854-1). Predicted N-terminus: Asp 24

#### **Molecular Characterization**

EDIL3(Asp 24 - Glu 480) Fc(Pro 100 - Lys 330)
O43854-1 P01857

This protein carries a human IgG1 Fc tag at the C-terminus.

The protein has a calculated MW of 77.8 kDa. The protein migrates as 80-90 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.

#### Endotoxin

Less than 1.0 EU per µg by the LAL method.

#### **Purity**

>95% as determined by SDS-PAGE.

#### **Formulation**

Lyophilized from  $0.22~\mu m$  filtered solution in 50~mM Tris, 100~mM Glycine, 25~mM Arginine, 150~mM NaCl, pH7.5 with trehalose as protectant.

Contact us for customized product form or formulation.

#### Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.

## **Storage**


For long term storage, the product should be stored at lyophilized state at -20  $^{\circ}\mathrm{C}$  or lower.

Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.

# **SDS-PAGE**




Human EDIL3, Fc Tag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95%.

## **Bioactivity-ELISA**








Biotinylated Human ITGAV&ITGB3 Heterodimer Protein, His, Avitag&Tag Free Conc. (μg/mL)

Immobilized Human EDIL3, Fc Tag (Cat. No. ED3-H5259) at 5  $\mu$ g/mL (100  $\mu$ L/well) can bind Biotinylated Human ITGAV&ITGB3 Heterodimer Protein, His,Avitag&Tag Free (Cat. No. IT3-H82W9) with a linear range of 0.02-0.625  $\mu$ g/mL (QC tested).

# **Bioactivity-SPR**



Human EDIL3, Fc Tag (Cat. No. ED3-H5259) immobilized on CM5 Chip can bind Human ITGAV&ITGB3 Heterodimer Protein, His Tag&Tag Free (Cat. No. IT3-H52E3) with an affinity constant of 38.9 nM as determined in a SPR assay (Biacore 8K) (Routinely tested).

# Human EDIL3, Fc Tag ELISA 0.1 μg of Biotinylated Human ITGAV&ITGB3 Heterodimer Protein, His,Avitag&Tag Free per well 3 2 EC50=10.96 ng/mL

Immobilized Biotinylated Human ITGAV&ITGB3 Heterodimer Protein, His,Avitag&Tag Free (Cat. No. IT3-H82W9) at 1  $\mu$ g/mL (100  $\mu$ L/well) on streptavidin (Cat. No. STN-N5116) precoated (0.5  $\mu$ g/well) plate can bind Human EDIL3, Fc Tag (Cat. No. ED3-H5259) with a linear range of 0.3-39 ng/mL (Routinely tested).

10

Human EDIL3, Fc Tag Conc. (ng/mL)

100

1,000

# **Background**

EDIL3, also known as Del1, is a novel extracellular matrix protein encoding three Notch-like epidermal growth factor repeats, an RGD motif, and two discoidin domains. EDIL3 is expressed in an endothelial cell-restricted pattern during early development. EDIL3 might involves autocrine angiogenic pathway for the embryonic endothelium, and this function is mediated in part by productive ligation of integrin alpha V beta 3. Overall, EDIL3 plays an important role in mediating angiogenesis and may be important in vessel wall remodeling and development. Moreover, It also influences endothelial cell behavior.

#### **Clinical and Translational Updates**

