

# PE-Labeled Monoclonal Anti-FMC63 Antibody, Mouse IgG1 (Y45) (Site-specific conjugation) (0.03% Proclin)

Catalog # FM3-PY54A2



BIOSYSTEMS  
**Acro**

## Source

PE-Labeled Monoclonal Anti-FMC63 Antibody, Mouse IgG1 is produced via site-specific conjugation of PE to Monoclonal Anti-FMC63 Antibody, Mouse IgG1 under optimal conditions with a proprietary technology.

*We carry another premium grade PE-Labeled Monoclonal Anti-FMC63 Antibody, Mouse IgG1 (Y45) (FM3-PY54G0), produced with the same production process but under more rigorous quality control system that incorporates a comprehensive set of tests including sterility and endotoxin tests. It is designed for cell isolation and cell culture applications in the early preclinical stage.*

## Application

Flow Cytometry (Evaluation of Anti-CD19 (FMC63 scFv) CAR Expression).  
Please note that this product is NOT compatible to streptavidin detection system.

## Clone

Y45

## Species

Mouse

## Isotype

Mouse IgG1/kappa

## Specificity

Specifically recognizes the antigen-recognition domain of FMC63 derived CARs.

## Immunogen

Recombinant FMC63 scFv derived from HEK293 cells.

## Conjugate

PE

Excitation Wavelength: 488 nm / 561 nm

Emission Wavelength: 575 nm

## Isotype Control

The Isotype control is sold separately and you can search for Cat. No. [DNP-PMI](#) for product information.

## Recommended Dilution

1:50

## Formulation

Lyophilized from 0.22 µm filtered solution in PBS, 0.5% BSA, 0.03% Proclin, pH7.4 with trehalose as protectant.

Contact us for customized product form or formulation.

## Reconstitution

Please see Certificate of Analysis for specific instructions.

*For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.*

## Storage

For long term storage, the product should be stored at lyophilized state at -20°C or lower.

*Please protect from light and avoid repeated freeze-thaw cycles.*

This product is stable after storage at:

- -20°C to -70°C for 24 months in lyophilized state;
- -70°C for 12 months after reconstitution;
- 2-8°C for 12 months after reconstitution.

## Bioactivity-FACS

Discounts, Gifts,  
and more!




➤ [www.acrobiosystems.com](http://www.acrobiosystems.com)


7/23/2024

# PE-Labeled Monoclonal Anti-FMC63 Antibody, Mouse IgG1 (Y45) (Site-specific conjugation) (0.03% Proclin)

Catalog # FM3-PY54A2



## 0.03% Proclin&Preservative free PE-Labeled Monoclonal Anti-FMC63 scFv Antibody, Mouse IgG1 FACS



5e5 of anti-CD19 CAR-293 cells were stained with 100  $\mu$ L of 1:50 dilution (2  $\mu$ L stock solution in 100  $\mu$ L FACS buffer) of PE-Labeled Monoclonal Anti-FMC63 Antibody, Mouse IgG1 (Cat. No. FM3-PY54A2) and isotype control antibody respectively. PE signal was used to evaluate the binding activity (QC tested).

Binding activity of the PE-Labeled Monoclonal Anti-FMC63 Antibody, Mouse IgG1 before and after adding 0.03% Proclin was evaluated in the above FACS analysis. The result shows that PE-Labeled Monoclonal Anti-FMC63 Antibody, Mouse IgG1 (Cat. No. FM3-PY54A2) and PE-Labeled Monoclonal Anti-FMC63 Antibody, Mouse IgG1 (Cat. No. FM3-HPY53) have the same binding activity against anti-CD19 CAR-293 cells.

## Background

FMC63 is an IgG2a mouse monoclonal antibody specific for CD19, which is a target for the immunotherapy of B lineage leukaemias and lymphomas. FMC63 scFv is the most commonly used ectodomain component of CD19-specific CARs. So far, most of reported CART19 trials contain the anti-CD19 scFv derived from FMC63, including the two FDA-approved CARs Kymriah and Yescarta.

## Clinical and Translational Updates

Discounts, Gifts,  
and more!

