

## Synonym

ERBB2,CD340,HER-2,neu,HER2,MLN19,NEU,NGL,TKR1

#### **Source**

Human Her2, His Tag(HE2-H5225) is expressed from human 293 cells (HEK293). It contains AA Thr 23 - Thr 652 (Accession # <u>P04626-1</u>). Predicted N-terminus: Thr 23

## **Molecular Characterization**

Her2(Thr 23 - Thr 652) P04626-1

Poly-his

This protein carries a polyhistidine tag at the C-terminus.

The protein has a calculated MW of 70.2 kDa. The protein migrates as 70-95 kDa when calibrated against <u>Star Ribbon Pre-stained Protein Marker</u> under reducing (R) condition (SDS-PAGE) due to glycosylation.

#### **Endotoxin**

Less than 0.2 EU per µg by the LAL method.

## **Purity**

>95% as determined by SDS-PAGE.

>90% as determined by SEC-MALS.

#### **Formulation**

Lyophilized from 0.22  $\mu m$  filtered solution in PBS, pH7.4 with trehalose as protectant.

Contact us for customized product form or formulation.

#### Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.

#### Storage


For long term storage, the product should be stored at lyophilized state at -20°C or lower.

Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 12 months under sterile conditions after reconstitution.


## **SDS-PAGE**



Human Her2, His Tag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95% (With <u>Star Ribbon Pre-stained Protein Marker</u>).

## **Bioactivity-ELISA**


## **SEC-MALS**




The purity of Human Her2, His Tag (Cat. No. HE2-H5225) is more than 90% and the molecular weight of this protein is around 75-105 kDa verified by SEC-MALS.

Report






Immobilized Human Her2, His Tag (Cat. No. HE2-H5225) at 1  $\mu$ g/mL (100  $\mu$ L/well) can bind Anti-Her2 antibody with a linear range of 0.061-1.95 ng/mL (QC tested).




Immobilized Human Her2, His Tag (Cat. No. HE2-H5225) at 1  $\mu$ g/mL (100  $\mu$ L/well) can bind Trastuzumab Biosimilar with a linear range of 0.1-2 ng/mL (Routinely tested).


## **Bioactivity-SPR**



Herceptin (Trastuzumab) captured on CM5 chip via anti-human IgG Fc antibodies surface, can bind Human Her2, His Tag (Cat. No. HE2-H5225) with an affinity constant of 1.07 nM as determined in a SPR assay (Biacore T200) (Routinely tested).



Immobilized Human Her2, His Tag (Cat. No. HE2-H5225) at 1  $\mu$ g/mL (100  $\mu$ L/well) can bind Pertuzumab Biosimilar with a linear range of 0.2-4 ng/mL (Routinely tested).




Immobilized Human Her2, His Tag (Cat. No. HE2-H5225) at 1  $\mu$ g/mL (100  $\mu$ L/well) can bind Bispecific Antibody (Her2 × Her3) with a linear range of 0.8-3 ng/mL (Routinely tested).





## **Bioactivity-BLI**



Loaded Herceptin (Trastuzumab) on AHC Biosensor, can bind Human Her2, His Tag (Cat. No. HE2-H5225) with an affinity constant of 0.825 nM as determined in BLI assay (ForteBio Octet Red96e) (Routinely tested).

# Background

Human Epidermal growth factor Receptor 2 (HER2) is also called ERBB2, HER-2,HER-2 /neu, NEU, NGL,TKR1 and c-erb B2, and is a protein giving higher aggressiveness in breast cancers. It is a member of the ErbB protein family, more commonly known as the epidermal growth factor receptor family. HER2 is a cell membrane surface-bound receptor tyrosine kinase and is normally involved in the signal transduction pathways leading to cell growth and differentiation. HER2 is thought to be an orphan receptor, with none of the EGF family of ligands able to activate it. Approximately 30% of breast cancers have an amplification of the HER2 gene or overexpression of its protein product. Overexpression of this receptor in breast cancer is associated with increased disease recurrence and worse prognosis. HER2 appears to play roles in development, cancer, communication at the neuromuscular junction and regulation of cell growth and differentiation.

# **Clinical and Translational Updates**

