

Synonym

Coagulation Factor III, Tissue Factor, TF, F3, CD142

Source

Mouse Tissue Factor, His Tag (TF3-M52H3) is expressed from human 293 cells (HEK293). It contains AA Ala 29 - Glu 251 (Accession # [P20352-1](#)).

Predicted N-terminus: Ala 29

Molecular Characterization

TF(Ala 29 - Glu 251) P20352-1	Poly-his
----------------------------------	----------

This protein carries a polyhistidine tag at the C-terminus.

The protein has a calculated MW of 27.3 kDa. The protein migrates as 40-55 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.

Endotoxin

Less than 1.0 EU per µg by the LAL method.

Purity

>90% as determined by SDS-PAGE.

>90% as determined by SEC-MALS.

Formulation

Lyophilized from 0.22 µm filtered solution in 50 mM Tris, 150 mM NaCl, pH7.5. Normally trehalose is added as protectant before lyophilization.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.

Storage


For long term storage, the product should be stored at lyophilized state at -20°C or lower.

Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:


- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.

SDS-PAGE



Mouse Tissue Factor, His Tag on SDS-PAGE under reducing (R) condition. The gel was stained overnight with Coomassie Blue. The purity of the protein is greater than 90%.

SEC-MALS



The purity of Mouse Tissue Factor, His Tag (Cat. No. TF3-M52H3) was more than 90% and the molecular weight of this protein is around 35-45 kDa verified by SEC-MALS.

[Report](#)

Background

Full-length tissue factor (TF) is a transmembrane receptor and cofactor for factor (F)VII/FVIIa. In addition to full-length TF, an alternative spliced (as) form of TF can be generated that lacks the transmembrane domain and is released from cells. In contrast to TF, asTF has low procoagulant activity because it lacks the transmembrane domain. Tissue factor is expressed by cells around blood vessels, such as adventitial fibroblasts, and body surfaces, such as epithelial cells, and plays

a critical role in hemostasis. TF also contributes to various forms of thrombosis. Many cancers, particularly adenocarcinomas, express high levels of TF. A high level of tumor TF expression is associated with poor prognosis in many types of cancers, including breast, prostate, colorectal, and pancreatic cancer.

Clinical and Translational Updates

Please contact us via TechSupport@acrobiosystems.com if you have any question on this product.