



Synonym

DIF,TNF-alpha,TNFA,TNFSF2,cachexin,cachectin,TNF α

Source

Biotinylated Human TNF-alpha Protein, Fc,Avitag(TNA-H82F3) is expressed from human 293 cells (HEK293). It contains AA Val 77 - Leu 233 (Accession # [P01375-1](#)).

Predicted N-terminus: Val 77

Molecular Characterization

TNF-alpha(Val 77 - Leu 233) P01375-1	Fc(Pro 100 - Lys 330) P01857	Avi
---	---------------------------------	-----

This protein carries a human IgG1 Fc tag at the C-terminus, followed by an Avi tag (Avitag™).

The protein has a calculated MW of 45.5 kDa. The protein migrates as 50-55 kDa when calibrated against [Star Ribbon Pre-stained Protein Marker](#) under reducing (R) condition (SDS-PAGE) due to glycosylation.

Labeling

Biotinylation of this product is performed using Avitag™ technology. Briefly, the single lysine residue in the Avitag is enzymatically labeled with biotin.

Protein Ratio

Passed as determined by the HABA assay / binding ELISA.

Endotoxin

Less than 1.0 EU per μ g by the LAL method.

Purity

>90% as determined by SDS-PAGE.

Formulation

Lyophilized from 0.22 μ m filtered solution in PBS, pH7.4 with trehalose as protectant.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.

Storage


For long term storage, the product should be stored at lyophilized state at -20°C or lower.

Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.


SDS-PAGE



Biotinylated Human TNF-alpha Protein, Fc,Avitag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 90% (With [Star Ribbon Pre-stained Protein Marker](#)).

Bioactivity-ELISA

SEC-MALS





The purity of Biotinylated Human TNF-alpha Protein, Fc,Avitag (Cat. No. TNA-H82F3) is more than 85% and the molecular weight of this protein is around 275-315 kDa verified by SEC-MALS.

[Report](#)


Discounts, Gifts,
and more!





Immobilized Human TNFR1, Fc Tag (Cat. No. TN1-H5251) at 5 µg/mL (100 µL/well) can bind Biotinylated Human TNF-alpha Protein, Fc,Avitag (Cat. No. TNA-H82F3) with a linear range of 0.1-3 ng/mL (QC tested).

Immobilized Human TNFR2, His Tag (Cat. No. TN2-H5227) at 5 µg/mL (100 µL/well) can bind Biotinylated Human TNF-alpha Protein, Fc,Avitag (Cat. No. TNA-H82F3) with a linear range of 0.1-6 ng/mL (Routinely tested).



Immobilized Humira at 1 µg/mL (100 µL/well) can bind Biotinylated Human TNF-alpha Protein, Fc,Avitag (Cat. No. TNA-H82F3) with a linear range of 0.1-6 ng/mL (Routinely tested).

Background

Tumor necrosis factor alpha (TNFα) is a cytokine produced primarily by monocytes and macrophages. It is found in synovial cells and macrophages in the tissues. The primary role of TNFα is in the regulation of immune cells. TNFα is able to induce apoptotic cell death, to induce inflammation, and to inhibit tumorigenesis and viral replication. Dysregulation of TNFα production has been implicated in a variety of human diseases, including major depression, Alzheimers disease and cancer. Recombinant TNFα is used as an immunostimulant under the INN tasonermin. TNFα can be produced ectopically in the setting of malignancy and parallels parathyroid hormone both in causing secondary hypercalcemia and in the cancers with which excessive production is associated.

Clinical and Translational Updates

